arXiv:1703.02369v1 [cond-mat.dis-nn] 7 Mar 2017

Applications of neural networks to the studies of phase transitions of two-dimensional Potts models

C.-D. Li,[1] D.-R. Tan,[1] and F.-J. Jiang[1,*]

[1]*Department of Physics, National Taiwan Normal University, 88, Sec.4, Ting-Chou Rd., Taipei 116, Taiwan*

We study the finite temperature (FT) phase transitions of two-dimensional (2D) q-states Potts models on the square lattice, using the first principles Monte Carlo (MC) simulations as well as the techniques of neural networks (NN). We demonstrate that the ideas from NN can be adopted to study these considered FT phase transitions efficiently. In particular, even with a simple NN constructed in this investigation, we are able to obtain the relevant information of the nature of these FT phase transitions, namely whether they are first order or second order. Our results strengthens the potential applicability of machine learning in studying various states of matters. Subtlety of applying NN techniques to investigate many-body systems is briefly discussed as well.

PACS numbers:

Introduction — While originally being a topic in computer science such as the artificial intelligence, the techniques of machine learning (ML) have been applied to many other fields of science in the last a few decades[1–8]. For example, the strategy of using methods of ML in analyzing large data sets of traffic flow and genome sequences has gained a lot of attention recently. Ideas from ML provide alternatives to the conventional approaches in various scientific areas. In some cases, the performance of these new approaches using the techniques of ML is as efficient as, or even better than the conventional ones. In the near future, it is anticipated that ML is likely to play a crucial role in many traditional fields of science.

The theoretical study of different phases of many-body systems, as well as the characters of the transitions between these various states is one of the major research topics in condensed matter physics[9]. Numerous numerical techniques, such as Monte Carlo simulations, series expansion, exact diagonalization, tensor networks and so on, are available for fulfilling these tasks[10,11]. Indeed, studies using these traditional methods have provided valuable information regarding the properties of matters. In particular, using the theoretical results obtained by these conventional techniques in conjunction with the data of relevant experiments, properties of some materials have been understood to an unprecedented precision.

With the advance of computing power as well as the development of ML algorithms, one interesting question arises, namely are these techniques of ML capable of efficiently studying different phases of matters? Such an idea has triggered pioneering works of applying ML algorithms to investigate phases, particularly the transitions between different states, of certain condensed matter systems[12–24]. In these pioneering studies, it is demonstrated that the neural networks (NN) of ML is a powerful approach to distinguish phases of many-body systems. Furthermore, NN can even detect exotic states such as a phase with topological order. Finally, NN is capable of locating the place in the relevant parameter space where the transition between two phases occurs as well.

Motivated by the success of applying NN to study phases of condensed matter systems, in this investigation we examine if one can obtain further information regarding a phase transition beyond detecting the location of the critical point. For instance, it will be extremely interesting to determine whether a studied phase transition is continuous or not, by employing the NN techniques. Remarkably, for the considered models here, namely the two-dimensional (2D) q-states Potts models with q being a positive integer[25–27], even a simple NN can provide us with relevant information to unambiguously determine the nature of the associated finite temperatures (FT) phase transitions of these models. Our investigation is not only important and interesting in itself, the obtained results also strengthen the applicability of NN in studying condensed matter systems.

Microscopic models, Observables, and Methods — The Hamiltonian H of the 2D q-states Potts model on the square lattice considered in our study is given by

$$\beta H = -\beta \sum_{\langle ij \rangle} \delta_{\sigma_i, \sigma_j}, \tag{1}$$

where β is the inverse temperature and $\langle ij \rangle$ stands for the nearest neighboring sites i and j. In addition, the Potts variable σ_i appearing above at each site i has the expression

$$\sigma_j = \exp\left(i\frac{2\pi s_j}{q}\right), s_j = 1, 2, 3, ..., q. \tag{2}$$

In the literature, the FT transitions of 2D q-states Potts models on the square lattice are second order for $q \leq 4$, and first order when $q \geq 5$. In this investigation, we have studied these FT phase transitions using both the method of Monte Carlo (MC) calculations and the techniques of NN. In particular, special attention has been focused on whether the methods of NN can be employed to determine the nature of these considered FT phase transitions efficiently.

To study these FT phase transitions through the MC simulations, the observable magnetization per site $\langle |m| \rangle$,

where m takes the following form

$$m = \frac{1}{L^2} \sum_i \sigma_i, \tag{3}$$

is measured. Here L is the linear box sizes used in the simulations. Besides, the energy density (energy per site) $\langle E \rangle$ is calculated as well. By studying the histograms of these two physical quantities, the nature of these FT phase transitions, i.e., first order or second order, can be effectively decided.

In addition to the conventional approach of using MC simulations, we also investigate these FT phase transitions with the techniques of NN. By considering quantities which are defined within the NN framework, we find that the nature of these FT phase transitions can be accurately determined as well. The procedures, including the considered quantities, associated with the constructed NN in this study will be described in details in the relevant sections.

The Numerical Results — To investigate the nature of the considered FT phase transitions, we have performed MC simulations for $q = 2, 3, 4, 5$, and 10, using the Swendsen Wang algorithm[28–30]. In particular, at every considered temperature close to the critical temperatures $T_c(q, L)$, a few thousand configurations are stored for each studied q and L. These configurations will be used in the NN approach to construct the appropriate quantities serving as the main information of deciding the nature of the considered FT phase transitions. In the next subsection, we will briefly present the numerical results obtained from the MC simulations.

Results of MC Simulations — One efficient method to determine the nature of a (FT) phase transition, namely to decide whether the transition is first order or second order, is to investigate the histograms of some relevant observables. Specifically, if a (FT) phase transition is first order, then for sufficiently large L, two peaks will appear in the histograms of certain observables when the simulations are carried out at (or close to) the corresponding critical point. Furthermore, the relative heights between the peak(s) and the trough (located in the middle) will approach infinity as one increases L. On the other hand, if the considered phase transition is second order, then the related histograms do not have these features. It should be pointed out that while for a second order phase transition, one may observe the appearance of two peaks in the histograms (of certain observables), the mentioned relative heights between the peak(s) and the trough will saturate to a constant. With this powerful idea, we indeed find that while the FT phase transitions for $q = 2, 3$, and 4 are continuous, the corresponding transitions of $q = 5$ and 10 are first order. Figures 1 and 2 are the histograms of $\langle |m| \rangle$ for 2- and 10-states Potts models, respectively (These data are determined with $L = 40$)[31]. Notice two peaks clearly show up in fig. 2 and the outcomes in that figure indicate that the FT phase transition for 10-states Potts model is first order. Furthermore, the results in fig. 1 are obtained at temperatures 1.133 (top) and 1.260 (bottom), and no two peaks structure is observed for other temperatures, separating from each other by 0.001, between $T = 1.133$ and $T = 1.260$. These two figures clearly show convincing signals that the FT phase transitions of 2- and 10-states Potts models are second order and first order, respectively. We have additionally simulated $L = 80$ lattices for 2-states Potts model and have found a similar scenario as that of $L = 40$. It should be pointed out that because $q = 4$ and $q = 5$ are at the edges of separating second and first order FT phase transitions of 2D Potts models on the square lattice, data of larger L are needed in order to reach a conclusive answer of whether the FT phase transitions of 4- and 5-states Potts models are second order or first order. Although our data for 4- and 5-states Potts model do not lead to absolutely conclusive results, they do show the correct trends.



FIG. 1: Histograms of $\langle |m| \rangle$ at $T = 1.133$ (top) and $T = 1.260$ (bottom) for 2D 2-states Potts model with $L = 40$ on the square lattice. Several ten thousand data, separating by few thousand updates from each other after the thermalization, are generated for each of the plots. The histograms are produced by the "hist" function of pylab[32].

Results of NN Methods — Besides the conventional approach, as shown in previous subsection, in the following we will employ the techniques of supervised ML, namely convolutional NN to study the considered FT phase transitions. In particular, the Keras library[33] is used in our investigation. The set up of the NN in this study consists of one input layer, one convolutional layer with kernel size 3 by 3, one average pooling layer with kernel size 2 by 2, and one output layer. The initial matrix elements W_{ij} connecting to the output layer, as well as the pooling kernel are the default ones of Keras. For a given fixed



FIG. 2: Histograms of $\langle|m|\rangle$ for $L = 20$ (top) and 40 (bottom) for 2D 10-states Potts model on the square lattice. Around twenty thousand data, separating by few thousand updates from each other after the thermalization, are generated for each of the plots. The histograms are produced by the "hist" function of pylab[32].

box size L and for each number m in $\{1, 2, 3, ..., q\}$, 200 identical L^2-vectors are generated. In particular, every element in these vectors has an initial numerical value m. These $200 \times q$ objects are then one-hot encoded in order to be used as the training set for the set up NN. Besides the investigation of using 200 L^2-vectors for each m in $\{1, 2, 3, ..., q\}$, we also consider the case of generating 400 L^2-vectors and find no significantly different results compared to those associated with the case of 200 L^2-vectors. The convolutional layer is activated by rectified linear functions. We have considered different initial filter kernels, and find that there are some restrictions on the corresponding kernel matrix elements. Specifically, each matrix element should not be too large, otherwise one hardly obtains satisfactory results. The main algorithm considered in the built NN is the mini-batch. Furthermore, we exploit the Adam optimizer of Keras to update the weights of the constructed NN. Various batch sizes, ranging from 10 to 80 are used. We find that the obtained results are stable with respect to the batch sizes as long as the used batch size is not (too) small or large. The number of epoch is confined by an early stop criterion and $L2$-regularization is imposed in calculating the loss functions (We use "categorical crossentropy" as the loss functions). Notice the output layer in the set up NN is activated by a softmax function and is a q-vector. The norms of these output vectors will be the main sources of deciding whether the studied FT transitions are first



FIG. 3: The set up NN in this study.



FIG. 4: Histograms of R for $L = 10$ (top) and 80 (bottom) for 2D 10-states Potts model. Around ten thousand data points close to the corresponding T_c are generated for each of the plots. Data are recorded once for every 2000 updates after the thermalization. The histograms are produced by the "hist" function of pylab[32].

order or second order. The pictorial representation of the set up NN is depicted in fig. 3. Notice there are several tunable parameters in our NN and we will make a remark regarding this later.

For each temperature, the testing set for the constructed NN are composed of several thousand configurations generated by MC simulations (separating from each other by few thousand updates). Notice on the one hand, the norms of the output q-vectors are 1 at temperatures below T_c. On the other hand, for temperatures above T_c, the corresponding output q-vectors should have the value of $\frac{1}{\sqrt{q}}$ as their norms. Based on this observation,

investigating the histograms of the norms R of the output q-vectors may shed some light on uncovering the nature of these FT phase transitions. For example, near T_c one may observe two peaks located at $R = 1$ and $R = 1/\sqrt{q}$. In particular, for a first order phase transition the relative heights between the peak(s) and the trough should increase with the linear box sizes L. Remarkably, such a scenario does happen in the histograms of R for 2D 10-states Potts model (See fig. 4). This implies that the FT phase transition of 2D 10-states Potts model is first order as expected. Notice the temperatures at which this two peaks phenomenon is the most obvious may be slightly different from those shown in fig. 2. This small discrepancy can be attributed to the fact that $T_c(L)$ are observable dependence. Figure 5 shows the histograms of R close to T_c for 2D 3-states Potts models. We find that as one moves from low temperatures to high temperatures, the histograms of R for 2D 3-states Potts model evolve from a configuration with (only) one peak locating near $R = 1$ to a configuration with (only) one peak near $R = \frac{1}{\sqrt{3}}$, without two peaks structure like those of figure 4 appearing for the intermediate temperatures. From this observation as well as the fact that in figure 5 the relative heights between the peaks and trough are almost the same for $L = 20$ and $L = 240$, one concludes that the corresponding FT phase transition is second order. These two results of 2D 3-and 10-states Potts models obtained using the constructed NN here agree with those determined by other traditional methods. Finally, by studying the histograms of R for 2D 2-,4-,and 5-states Potts models, we arrive at conclusions consistent with the known results in the literature regarding the nature of these FT phase transitions as well.



FIG. 5: Histograms of R for $L = 20$ (top) and 240 (bottom) for 2D 3-states Potts model. Around ten thousand data close to the corresponding T_c are generated for each of the plots. Data are recorded once for every 2000 updates after the thermalization. The histograms are produced by the "hist" function of pylab[32].

Discussions and Conclusions — In this study, we investigate the FT phase transitions of various 2D q-states Potts models using both the MC simulations and the techniques of NN. Special attention has been focused on whether the NN methods can be applied efficiently to determine the nature of the considered FT phase transitions. Remarkably, while one expects MC simulations is capable of carrying out such tasks, it is surprising that the simple NN constructed here provides us with clear signals to decide whether these FT phase transitions are first order or second order. Many pioneering works have demonstrated that NN is a powerful tool to distinguish phases of matters, and the results shown here certainly strengthen the applicability of NN in investigating condensed matter systems. One natural next step will be to build a NN capable of accurately calculating physical quantities, such as the critical exponents, of second order phase transitions.

In computer science like the artificial intelligence, the ideas behind designing algorithms are sophisticated so that the constructed algorithms can apply for computations in many different areas. Such a concept is well-suited in majority of scientific fields including physics. Notice different physical systems have their own special characters. Hence constructing appropriate NN based on certain properties of the considered physical systems is also an efficient way to conduct the investigation. In particular, several professional ML packages and libraries like Keras and TensorFlow are publicly available[33–36]. These ML packages and libraries can be easily adopted to study specific systems. Hence for systems with less complexity, the strategy considered here, namely designing the NN according to the characters of the investigated models is an effective approach as well. Finally, we would like to point out that a typical NN has several tunable parameters. For example, for the constructed NN here, the batch-size, the size of convolution kernel, as well as the number of training objects are all adjustable variables. For the obtained results to be reliable, one has to make sure that the conclusions stay the same when the tunable parameters are reasonably varied. For instance, we have tested the results determined by using filter kernels with different sizes and matrix elements. In addition, We have also considered the situations of modifying the average pooling layer. Based on these new analysis we find that while in the case of $L = 240$ one may not be able to arrive at satisfactory outcomes, the resulting histograms of R greatly favor the conclusions as those presented in previous sections regarding the nature of these considered FT phase transitions, provides that the tunable parameters are reasonably varied[37].

Partial support from Ministry of Science and Technology of Taiwan is acknowledged. We thank D. Banerjee

for reading the manuscript carefully and giving us very useful comments.

* fjjiang@ntnu.edu.tw

[1] Trevor Hastie, Robert Tibshirani, and Jerome Friedman, *The Elements of Statistical Learning: Data Mining, Inference, and Prediction, Second Edition* (Springer, 2009).

[2] Steven Horvath, *Weighted Network Analysis: Applications in Genomics and Systems Biology* (Springer Science+Business Media, LLC 2011).

[3] Laszlo Gyorfi, Gyorgy Ottucsak, and Harro Walk, *Machine Learning For Financial Engineering* (Imperial College Press, 2012).

[4] M. Kircher, Martin Kircher, Daniela M Witten, Preti Jain, Brian J O'Roak, Gregory M Cooper, and Jay Shendure, Nature Genet. **46**, 310315 (2014).

[5] John D. Kelleher, Brian Mac Namee, and Aoife D'Arcy, *Fundamentals of Machine Learning for Predictive Data Analytics: Algorithms, Worked Examples, and Case Studies* (MIT Press, 2015).

[6] Maxwell W. Libbrecht and William Stafford Noble, Nature Reviews Genetics **16**, 321332 (2015).

[7] Ian Goodfellow and Yoshua Bengio, *Deep Learning (Adaptive Computation and Machine Learning series)* (MIT Press, 2016).

[8] Adalbert F.X. Wilhelm and Hans A. Kestler, *Analysis of Large and Complex Data (Studies in Classification, Data Analysis, and Knowledge Organization)* (Springer International Publishing Switzerland, 2016).

[9] S. Sachdev, *Quantum Phase Transitions* (Cambridge University Press, Cambridge, 1999).

[10] A. W. Sandvik, *AIP Conference Proceedings* **1297** (2010).

[11] A. Avella and F. Mancini, *Strongly correlated Systems: Numerical Methods* (Springer Berlin Heidelberg, 2013).

[12] Juan Carrasquilla, Roger G. Melko, arXiv:1605.01735.

[13] Giuseppe Carleo, Matthias Troyer, arXiv:1606.02318.

[14] Giacomo Torlai and Roger G. Melko, Phys. Rev. B **94**, 165134 (2016).

[15] Peter Broecker, Juan Carrasquilla, Roger G. Melko, and Simon Trebst, arXiv:1608.07848.

[16] Kelvin Ch'ng, Juan Carrasquilla, Roger G. Melko, and Ehsan Khatami, arXiv:1609.02552.

[17] Akinori Tanaka, Akio Tomiya, arXiv:1609.09087.

[18] Evert P.L. van Nieuwenburg, Ye-Hua Liu, Sebastian D. Huber, arXiv:1610.02048.

[19] Junwei Liu, Huitao Shen, Yang Qi, Zi Yang Meng, Liang Fu, arXiv:1611.09364.

[20] Xiao Yan Xu, Yang Qi, Junwei Liu, Liang Fu, Zi Yang Meng, arXiv:1612.03804.

[21] Lei Wang, Phys. Rev. B **94**, 195105 (2016)

[22] Li Huang and Lei Wang, Phys. Rev. B **95**, 035105 (2017).

[23] Junwei Liu, Yang Qi, Zi Yang Meng, Liang Fu, Phys. Rev. B **95**, 041101 (2017).

[24] Qianshi Wei, Roger G. Melko, Jeff Z. Y. Chen, arXiv:1701.04390.

[25] K. Binder, Journal of Statistical Physics 24(1), 69-86 (1981).

[26] F. Y. Wu, Rev. Mod. Phys. **54**, 235 (1982).

[27] Alain Billoire, arXiv:hep-lat/9501003.

[28] R. H. Swendsen, and J.-S. Wang, (1987), Phys. Rev. Lett. **58(2)**, 86 (1987).

[29] M. E. Newman and G. T. Barkema, *Monte Carlo Methods in Statistical Physics* (Clarendon Press, 1999).

[30] David P. Landau and Kurt Binder, *A Guide to Monte Carlo Simulations in Statistical Physics* (Cambridge University Press, 2014).

[31] In this study, we do not carry out a detailed error analysis since very fine histograms of the relevant observables are used for the investigation.

[32] All the histograms shown in this study are produced by the "hist" function of pylab. This function has an input parameter n which can be a positive integer. The positive integer n is used to generate histogram of n-bins, including empty ones, of rectangular shape with equal width. For better visualization, each histograms presented here may be obtained using different number of bins.

[33] https://keras.io

[34] http://scikit-learn.org/stable/

[35] https://www.tensorflow.org

[36] http://deeplearning.net/software/theano/

[37] While the conclusions regarding the nature of the considered FT phase transitions are unchanged, the corresponding $T_c(q, L)$ at finite L may slightly vary when the used NN is modified. This effect can be treated as one part of the errors for $T_c(q, L)$ related to the NN methods.